

**ACTION PLAN FOR THE CONSERVATION OF
THE CORALLIGENOUS AND OTHER CALCAREOUS
BIO-CONCRETIONS IN THE MEDITERRANEAN SEA**



United Nations
Environment Programme



Mediterranean Action Plan
Barcelona Convention

Graphic layout and production

Final layout and production were prepared by the Specially Protected Areas Regional Activity Centre (SPA/RAC), Tunis.



Specially Protected Areas
Regional Activity Centre

Cover photos

SPA/RAC, University of Seville

UN Environment/Mediterranean Action Plan Barcelona Convention Secretariat

48, Vassileos Konstantinou Avenue 11635
P.O. Box: 18019, Athens Greece
Tel: +302107273100 - Fax: +30 2107253196
www.unepmap.org

Legal Notice

The designations employed and the presentation of the material in this document do not imply the expression of any opinion whatsoever on the part of UN Environment/Mediterranean Action Plan (MAP) concerning the legal status of any State, Territory, city or area, or of its authorities, or concerning the delimitation of their frontiers or boundaries.

Copyright

This publication may be reproduced in whole or in part and in any form for educational or non-profit purposes without special permission from the copyright holder, provided acknowledgement of the source is made. UN Environment/Mediterranean Action Plan would appreciate receiving a copy of any publication that uses this publication as a source. This publication cannot be used for resale or for any other commercial purpose whatsoever without permission in writing from UN Environment/Mediterranean Action Plan.

© 2017 United Nations Environment Programme / Mediterranean Action Plan
P.O. Box 18019, Athens, Greece

For bibliographic purposes this volume may be cited as:

Action Plan for the Conservation of the Coralligenous and Other Calcareous Bio-concretions in the Mediterranean Sea.
UN Environment/MAP Athens, Greece 2017.

TABLE OF CONTENTS

Decision IG.22/12	4
I. Current Situation Of Coralligenous Assemblages	5
I.1. Current knowledge	5
I.2. Distribution	5
I.3. Composition	5
II. Data collection and inventories	7
II.1. Specific inventories	7
II.2. Sites of particular interest	7
III. Monitoring activities	8
III.1. Types of monitoring	8
III.2. Monitoring methods	9
III.2.1. General considerations for sampling strategies for monitoring schemes	9
III.2.2. Spatial scales	9
III.2.3. Temporal scales	9
III.2.4. Sampling techniques	9
IV. Research activities	10
IV.1. Taxonomy	10
IV.2. Long term evolution	11
IV.3. Functioning	11
V. Conservation activities	11
V.1. Major threats	11
V.1.1. Trawling	12
V.1.2. Artisanal and recreational fishing	12
V.1.3. Anchoring	12
V.1.4. INVASIVE SPECIES	12
V.1.5. Global warming	12
V.1.6. Waste water discharges	13
V.1.7. Aquaculture	13
V.1.8. Changes in land use and coastal infrastructure construction and urbanization	13
V.1.9. Recreational activities (excluding fishing)	13
V.1.10. Mucilaginous and filamentous algal aggregates	13
V.2. Legislation and regulations	13
V.3. Creation of marine protected areas	14
V.4. National plans	14
VI. Coordination of this Action plan with other tools and initiatives	15
VII. Regional coordination structure	15
VIII. Participation in the implementation	15
Annex: implementation timetable	16
IX. References	17

Decision IG.22/12

Updated Action Plans Concerning “Cetaceans”, “Coralligenous and Other Calcareous Bio-concretions”, and “Species Introductions and Invasive Species”; Mandate for update of the “Action Plan on Marine and Coastal Birds” and revision of the “Reference List of Marine and Coastal Habitat Types in the Mediterranean”

The 19th Meeting of the Contracting Parties to the Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean, hereinafter referred to as “the Barcelona Convention”,

Recalling Articles 11 and 12 of the Protocol concerning Specially Protected Areas and Biological Diversity in the Mediterranean, hereinafter referred to as the “SPA/BD Protocol”, on national measures for the protection and the conservation of species and on the formulation and implementation of action plans for their conservation and recovery respectively;

Recalling Decision IG.19/12 of COP 16 (Marrakesh, Morocco, November 2009) related to the Amendments of the list of Annexes II and III of the SPA/BD Protocol and more particularly the marine and coastal bird species included then in Annex II to the SPA/BD Protocol “List of endangered or threatened species”;

Recalling Decision IG.20/4 and IG.21/3 of COP 17 (Paris, France, February 2012) and COP 18 (Istanbul, Turkey, December 2013) respectively adopting Ecological Objectives, Operational Objectives, GES and related targets ;

Recalling Decision IG.21/17 on the Programme of Work of COP 18 (Istanbul, Turkey, December 2013) on the update of the Action Plan for the Conservation of Cetaceans in the Mediterranean Sea and the Action Plan for the Conservation of the Coralligenous and Other Calcareous Bio-concretions in the Mediterranean Sea;

Having considered the report of the 12th Meeting of SPA/RAC Focal Points (Athens, Greece, May 2015);

1. **Adopts** the Updated Action Plan for the Conservation of Cetaceans in the Mediterranean Sea, as contained in Annex I to this Decision;
2. **Adopts** the Updated Action Plan for the Conservation of the Coralligenous and Other Calcareous Bio-concretions in the Mediterranean Sea, as contained in Annex II to this Decision;
3. **Adopts** the Updated Action Plan concerning Species Introductions and Invasive Species in the Mediterranean Sea, as contained in Annex III to this Decision;
4. **Requests** the Contracting Parties to take the necessary measures for the implementation of the updated Action Plans and report according to the cycle and format of the MAP/Barcelona Convention reporting system;
5. **Requests** SPA/RAC to provide support to the full implementation of the updated Action Plans;
6. **Requests also** SPA/RAC to update the Action Plan for the Conservation of Bird Species listed in Annex II to the SPA/BD Protocol including all 25 target species, and to revise the Reference List of Marine and Coastal Habitat Types in the Mediterranean for consideration by COP 20, taking in full account the biodiversity-related MAP Ecological Objectives, IMAP, and GES targets.

I. CURRENT SITUATION OF CORALLIGENOUS ASSEMBLAGES

I.1. CURRENT KNOWLEDGE

1. At present there is a general knowledge on the distribution, species composition and functioning of coralligenous assemblages and other calcareous bio-concretions. However, and despite the efforts conducted since the adoption in 2008 of the Action Plan for Coralligenous and other Calcareous Bioconstructions, in the Mediterranean, there are essential questions that need to be addressed to guarantee the conservation of these emblematic Mediterranean habitats (see specific sections).
2. Probably the number and quality of presentations during the 2nd Mediterranean Symposium on the Conservation of coralligenous and other calcareous bio-concretions (Portorož, Slovenia 29-30 October 2014) are the best example on the interest of Mediterranean scientific/managers community to improve the knowledge on these assemblages (Proceedings 2nd MSC&CBC 2014).
3. Despite of this, it was also noted that
 - (i) most actions regarded individual- national- based efforts and
 - (ii) the lack of structures for coordination in an efficient way of regional and/or pan-Mediterranean research actions.

There was a general consensus at the Symposium to establish a series of Working Groups to coordinate the human and resources in order to provide the needed general view on the coralligenous/maërl assemblages and overcome these gaps.

I.2. DISTRIBUTION

4. Coralligenous buildups and maërl beds are common all around the Mediterranean coasts, even in the easternmost coasts (Giakoumi et al. 2013, Martin et al. 2014). The coralligenous habitats are abundant in the Adriatic, Aegean and Tyrrhenian Seas as well as in the Algero-Provençal Basin. The coralligenous is less abundant in the Levantine Sea and Tunisian Plateau/Gulf of Sidra (Martin et al. 2014). Overall, data available cover approximately 30 % of the Mediterranean coasts while for the remaining 70 % no information was found (Martin et al. 2014). Regarding the depth distribution, most information concern the 10 to 50 m depth less information exists for the deeper range of distribution of coralligenous 50-200 m depth. Besides these large-scale assessments on distribution, at local scale some progress in cartographical data have been acquired in some areas especially in marine protected areas (e.g. Réserve Naturelle de Scandola, Parc National de Zembra, Area Marina Protetta di Tavolara Punta Coda Cavallo, Zakynthos Marine Protected Area). Overall, we lack of a complete and precise distribution information on coralligenous and other calcareous bioconstructions habitats.
5. The main constraints to provide a global view on the distribution of coralligenous and other calcareous habitats are the 1) their intrinsic heterogeneous distribution related with the spatial patterns of the geophysical and oceanographic conditions allowing their development, and 2) the technical and financial constraints of field mapping operations resulting in an unbalanced mapping efforts across the Mediterranean.
6. Geographical as well as depth distributional data are essential in order to know the real extent of these assemblages in the Mediterranean Sea as well as to implement appropriate management measures to guarantee their conservation.

I.3. COMPOSITION

7. Coralligenous concretions are the result of the building activities of algal and animal builders and the physical as well as biological eroding processes. The final result is a very complex structure composed of several microhabitats. Environmental factors (i.e., light, water movement and sedimentation rates) can vary by orders of magnitude in parts of the same concretion situated very close to each other. This great environmental heterogeneity allows several different assemblages to coexist in a reduced space. Assemblages situated in open waters (from horizontal to almost vertical surfaces) can be easily distinguished from those situated in overhangs and cavities.

8. Algae usually dominate in horizontal to sub-horizontal surfaces although their abundance decreases with decreasing irradiance. Two main algal communities have been distinguished in the western Mediterranean: an assemblage dominated by *Halimeda tuna* and *Mesophyllum alternans* (*Lithophyllo-Halimedetum tunae*), thriving in relatively high light levels, and an assemblage dominated by encrusting corallines (*Lithophyllum frondosum*, *L. cabiochae*, *Neogoniolithon mamillosum*) and *Peyssonnelia rosamarina* (*Rodriguezelletum strafforelloi*), and receiving low light levels.
9. Animal assemblages can greatly differ according to light levels reaching the coralligenous outcrop but also in relation to current intensity, sedimentation rates and geographical areas. In the richest, relatively more eutrophic zones, with rather constant and low water temperature, gorgonians usually dominate the community, but they are completely absent or rare in the more oligotrophic or low-current areas with rather high or seasonally variable temperature, being replaced by sponges, bryozoans or ascidians.
10. Maërl beds are also very diverse. Even if corallines are the main constituents (*Spongites fruticulosus*, *Lithothamnion corallioides*, *Phymatolithon calcareum*, *Lithothamnion valens*, *Lithothamnion minervae*, *Lithophyllum racemus*, *Lithophyllum frondosum*, and others), *Peyssonnelia* species (mainly *Peyssonnelia rosamarina*) can also be very important. The cover of erect algae depends on each particular site, displaying several facies (*Osmundaria volubilis*, *Phyllophora crispa*, *Kallymeniales*, *Laminaria rodriguezii*).
11. The group of experts in Tabarka suggested using the Reference List of Habitat types appearing in the Standard Data-entry Form (SDF) for National Inventories when looking for the composition of coralligenous assemblages. In 2011 a list of species to be considered in the inventory and/or monitoring of coralligenous communities was provided by UNEP-MAP-SPA/RAC (2011)¹. The species were arranged in the following categories:
 - Algal builders
 - Animal builders
 - Agglomerative' animals
 - Bioeroders
 - Species of particular importance (particularly abundant, sensitive, architecturally important or economically valuable)
 - Invasive species
12. The characterization of coralligenous based on the above-mentioned categories list can greatly help in our understanding on the coralligenous patterns across the Mediterranean. Since different regions and areas within regions are characterized by different composition, the assessment considering the proposed morpho-functional categories can provide an interesting comparative basis towards a general view on Mediterranean coralligenous assemblages. This approach besides the composition data would provide a functional perspective which greatly facilitate the development of indicators for the monitoring of the Good Environmental Status (GES) within the Marine Strategy Framework Directive and "COP18 EcAp Decision" (see Legislation and regulation section).
13. The suggestion when describing the composition of the coralligenous assemblages or the maërl beds would be to provide quantitative or semi-quantitative estimate on the abundance of typical/indicator species. Different visual and photographic methods as well as the combination of both have been proposed to obtain abundance estimates. For instance, the adoption of Braun-Blanquet (1979) methodology for marine assemblages characterization (Cebrian & Ballesteros, 2004). Through these assessments besides composition data, the abundance estimates of species found in the considered categories would provide insights in the ecological/conservation status of assemblages. For instance, the presence of invasive species (either alien or not normally occurring in the habitat) are often considered very good indicators of poor conservation status.
14. For maërl beds assemblages the same approach could be addressed although the current knowledge need to be improved to better define the categories and composition lists. In maërl beds, description is also possible naming the main maërl species and erect algae, as well as the main macroinvertebrates.

¹ Proposal of standard methods for inventorying and monitoring coralligenous and rhodoliths populations UNEP-MAP-SPA/RAC (2011)

II. DATA COLLECTION AND INVENTORIES

II.1. SPECIFIC INVENTORIES

15. As mentioned the coralligenous habitat includes several assemblages due to its great heterogeneity. There is a small scale heterogeneity in environmental factors throughout the coralligenous outcrops that determine different micro-habitats containing different species. In the surface of coralligenous outcrops, coralline algae usually dominate, together with a variable amount of erect algae and of suspension-feeders. Holes and cavities within the coralligenous structure sustain complex communities without algae and dominated by suspension-feeders. Small crevices and interstices are inhabited by a diverse endofauna, while many vagile species swarm everywhere, thriving also in the small patches of sediment retained by the framework. Large fishes (e.g. *Epinephelus marginatus*, *Scorpaena scrofa*, *Phycis phycis*) and decapods (e.g. *Palinurus elephas*, *Homarus gammarus*) dwell in the coralligenous assemblages. One of the consequences of this great environmental heterogeneity is the presence of a high biodiversity and a wide array of organisms in each coralligenous outcrops.
16. Maërl beds are considerably less complex than coralligenous outcrops although they have some epiflora and epifauna that are more related to plants and animals usually found in rocky substrata, but also they harbour typically invertebrates from sedimentary bottoms.
17. A considerable amount of research has been done on the biodiversity hosted by coralligenous outcrops. Ballesteros (2006) estimates a preliminary account of up to 1666 species at the scale of the Mediterranean Sea. However these estimates are far from providing us a general view of biodiversity dwelling in the coralligenous assemblages. There are at least two levels of information which should be considered i) in fine detailed taxonomic studies specially in less studied groups and ii) comprehensive biodiversity surveys in targeted geographical areas. This information would be complemented by the determination of typical/indicator species of coralligenous in different areas/regions across the Mediterranean (see Point 1.3. Composition).
18. Overall with this information we could improve the estimates on the total number of species associated to the coralligenous and analyze geographical variability biodiversity patterns considering different spatial scales. The same approach should be adopted for maërl beds.
19. Methods: For data collection several methodologies have been used in sampling rocky benthic systems and maërl beds (e.g. Bianchi et al., 2004, Kipson et al. 2011, Cechi et al. 2010, Gatti et al. 2015) and all of them present advantages and disadvantages. Moreover, suitability of each sampling method depends on the purposes of the study and on the taxonomic group considered.
20. As no sampling methodology can be universally applied, a general recommendation when making the assessments on species composition is to take into account the following considerations:
 - Use quantitative or semi-quantitative surveys instead of qualitative surveys wherever possible.
 - Clearly state the sampling and quantification methodology, including the period of the year, in order that it could be repeated in the future by independent teams for further comparison of data.
 - Samples have to be geographically positioned in the most accurate way.
 - Sampling has to be representative. Therefore, sampling areas should be larger than minimal sampling areas. It has to be noted that different taxonomic groups must be sampled using completely different representative areas.
 - Use photographic surveys to help in the identification of species.

II.2. SITES OF PARTICULAR INTEREST

21. Since the coralligenous and maërl assemblage in general thrive in deep waters, it is difficult to have an appropriate coverage of all the entire distribution range of the assemblage. Thus, it is recommended that inventories and monitoring be performed in selected sites of particular interest. The sites selection should be based in the most accurate previous information on the distribution, extension and ecological features and conservation status of coralligenous and maërl assemblages.

22. Amongst the criteria to be used in this selection, it is recommended the following ones:

- Existence of previous information on coralligenous assemblages or maërl beds at the site or, if there is no available information at all, sea bottom geomorphological features suitable for the development of coralligenous frameworks and/or rhodolites.
- Representativity of the coralligenous assemblages/maërl beds at a wide geographical area, whenever it is possible, according to present knowledge.
- Existence of control and/or management of anthropic activities at the site. In this sense, marine protected areas are suitable places to be selected.
- Especially healthy coralligenous and maërl assemblages are worth to be selected in order to assess the reference conditions.
- Coralligenous communities and maërl beds under the effects of direct or indirect anthropogenic disturbances are worth to be selected in order to assess the impact conditions.

III. MONITORING ACTIVITIES

23. Even if coralligenous/maërl assemblages are characterized by very slow dynamics (Garrabou et al., 2002, Teixidó et al. 2011), at least in the absence of punctual catastrophic disturbances (Teixidó et al. 2013), developing monitoring activities is of great interest to track their conservation status and detect changes associated to pressure and punctual human- as well as natural processes-related disturbances .
24. Monitoring is necessary to understand the processes behind long-term dynamics in the assemblages and is a central element for the implementation and evaluation of efficient management plans. Besides, the monitoring activities on coralligenous assemblages are required for the implementation of European Marine Strategy Framework Directive (MSFD 2008/56/EC) and the Barcelona Convention Decision² (see Legislation and regulation section) seeking to maintain the Good Environmental Status of assemblages.

III.1. TYPES OF MONITORING

25. The basic scheme of surveillance includes periodic monitoring of reference parameters (indicators) informing about the conservation status of coralligenous/maërl assemblages. The monitoring should be designed to be as simple as possible. Neither standard methods have been proposed nor environmental nor ecological quality indexes have been established yet for the coralligenous assemblages.
26. Due to the heterogeneity and habitat complexity monitoring should be conducted by a combination of methods to gather habitat, species and degree of impacts data.
27. Monitoring parameters should provide information on:
- Structural and functional parameters of assemblages:
- Species/Categories composition/abundance (semi- or quantitative data)
 - Indicators on the degree of complexity of coralligenous habitats
 - Indicators on coralligenous functioning: bioeroders and bioconstructors
 - Qualitative, semi- and quantitative indicators on the impacts of different disturbances on coralligenous communities (e.g. presence of fishing nets, invasive species, high diving pressure)
- Environmental parameters
- Temperature, sedimentation.

² Decision IG.21/3 on the Ecosystems Approach including adopting definitions of Good Environmental Status (GES) and targets

III.2. MONITORING METHODS

III.2.1. GENERAL CONSIDERATIONS FOR SAMPLING STRATEGIES FOR MONITORING SCHEMES

28. Bearing in mind the depth distribution of coralligenous/maerl assemblages monitoring methods have to be adapted to limited bottom working time by scuba divers (due to long decompression times and limitation of diver performance in deep waters; Tetzaff & Thorsen, 2005; Germonpre, 2006) and the limitation of the use of Remote Operate Vehicles (ROVs) beyond the operational depth of scuba divers (0-40 m).

III.2.2. SPATIAL SCALES

29. The high scale heterogeneity of coralligenous outcrops implies a large sampling area to be representative (Ballesteros, 2006). At present, some studies have determined the minimum sampling areas in some assemblages (Kipson et al. 2011), similar approaches should be carried out in other coralligenous morpho-types. In general, in order to gather relevant data on the different indicators in each monitoring site the total sampling area (including different replication strategies) should cover about 5 to 30 m² (Deter et al. 2012, Garrabou et al. 2014, Gatti et al. 2015).
30. At each site, determine a specific depth range where the monitoring will be carried out (e.g. 30-35 m), in order to avoid the potential effect of depth on the outcome of the surveys. Within the depth range selected, in order to limit the effects of local heterogeneity on the outcome of the surveys, determine when possible, with the help of remarkable seascape marks, the specific monitoring area (e.g. it should be an area of several 100 m²) of each sampling site. Eventually some marks can be fixed to help the sample in the same monitoring area. Finally, in each targeted geographic areas several sites should be monitored in order to better infer the conservation trends of assemblages.
31. When selecting monitoring sites one should keep in mind the existence of previous information on the extension and ecological quality of the coralligenous habitat. During selection process, it is recommended to consider the following questions:
- Is there previous information available on coralligenous assemblages at the site or, if there is no available information at all, are the sea bottom geomorphological features suitable for the development of coralligenous frameworks?
 - According to the present knowledge, are the considered coralligenous assemblages representative for a wider geographical area?
 - Are the considered coralligenous assemblages especially healthy to be able to serve as reference points?
 - Are the considered coralligenous assemblages under some clearly recognizable direct or indirect anthropogenic disturbance that would allow the assessment of the impact of these disturbances?

III.2.3. TEMPORAL SCALES

32. The low dynamic of coralligenous assemblages (Garrabou et al., 2002, Casas et al. 2015) allows to set the sampling periodicity between 3-5 years for monitoring purposes. Regarding the period of monitoring, the ideal period is late summer (late August to early October). At that time water transparency and temperature allow better performances on data gathering and photosampling. In addition, if any mass mortality occurred during summer it can be observed in this period.

III.2.4. SAMPLING TECHNIQUES

33. During the last years different approaches have been adopted for the assessment of conservation status of coralligenous assemblages using visual and/or photographic surveys (e.g. Cormaci et al., 2004, Kipson et al. 2011, Deter et al. 2012, Garrabou et al. 2014, Gatti et al. 2015). The sampling approaches developed are based on non-destructive methods aiming to furnish rapid quantitative and semi-quantitative assessments of different parameters.

34. The basic parameters assessed by photographic sampling and visual census are abundance (e.g. coverage, density) of species found in the assemblages and estimations on the degree of impact of different key processes (e.g. mortality events, bioerosion, fishing) related with the conservation of coralligenous assemblages.
35. Monitoring of environmental parameters is also needed if we want to relate changes in the coralligenous/maërl assemblages with disturbances related to hydrographic conditions. The most important variables to be monitored are: water temperature, sedimentation rates, nutrient concentration in seawater, particulate organic matter and water transparency.
36. Different initiatives (this Action Plan and EU directives) are focused on the development of indicators about the conservation and good environmental status of coralligenous. Through the monitoring activities presented we could obtain useful indicators (See Annex). These indicators are intended to inform decision makers and stakeholders and to support conservation and management planning (including MPAs network design) to guarantee the conservation of the coralligenous habitat.
37. Standardized protocols for the characterization of coralligenous/maërl assemblages need to be developed. The main goal of this action would be to do a comparative evaluation of the tools and sampling designs to be applied for the characterization of coralligenous habitats (e.g. in terms of species diversity (α , β , γ), structural complexity and main ecological processes) and to assess the level of impact of human pressures.
38. Indices and/or intercalibration initiatives to determine environmental status of coralligenous conservation should be developed to analyze the available indices developed to determine the Good Environmental Status of coralligenous to provide a common framework to compare the status of coralligenous across the Mediterranean.

IV. RESEARCH ACTIVITIES

IV.1. TAXONOMY

39. Coralligenous/maërl assemblages probably are two of the most important hot-spots of species diversity in the Mediterranean, together with *Posidonia oceanica* meadows (Ballesteros, 2006; BIOMAERL team, 2003). In comparison to the large amount of literature devoted to the study of *Posidonia oceanica* meadows, studies devoted to strengthen the knowledge of coralligenous/maërl biodiversity are scarce. Therefore, due to the rich fauna, high heterogeneity at all scales, and complex structure of coralligenous/maërl assemblages, together with the paucity of studies dealing with coralligenous/maërl biodiversity, it can be assumed that at least coralligenous assemblages harbour more species than any other Mediterranean community. The check-list proposed in the second chapter of this Action plan should cover all the species found to date in coralligenous/maërl communities. However research in taxonomy is also needed as a large amount of taxonomic groups absolutely lack not only of a comprehensive study but almost any study dealing with species which can be found in coralligenous outcrops or maërl beds. The use of genetic tools can help in resolving taxonomic “problems” and discovering cryptic species (e.g. Dailianis et al. 2014).
40. Taking into account the current knowledge of biodiversity in coralligenous/maërl communities (Ballesteros, 2006), the following taxonomic groups need an important investment in research:
 - Copepods
 - Cumaceans
 - Isopods
 - Molluscs
 - Mysids
 - Nematods
 - Nemertean
 - Ostracods
 - Phyllocarids
 - Polychaeta
 - Pycnogonids
 - Tanaidaceans

41. Further research on other groups is also acknowledged, as it will surely provide new reports of species for coralligenous outcrops and maërl beds.

IV.2. LONG TERM EVOLUTION

42. To understand long-term dynamics of coralligenous assemblages in some selected areas sentinel/reference sites should be setup. Processes taking place in coralligenous communities in absence of disturbances usually display slow dynamics – i.e. decades – (Garrabou *et al.*, 2002). Population dynamics of outstanding and key species show low growth rates and low population dynamics (e.g. Coma *et al.* 1998, Teixidó *et al.* 2011). Therefore, even if some of the patterns and processes that have been described so far occur in short time periods (e.g. mortality events; Cerrano *et al.*, 2000; Garrabou *et al.*, 2009), evolution of coralligenous can only be understood from a long-term perspective. Maërl beds are even less known as there are no comprehensive revisions in this subject regarding Mediterranean rhodolits.
43. Sentinel/reference sites are recommended to be visited once a year to obtain a robust temporal series. Even if seasonality in coralligenous/maërl communities is not as important as it is in shallower environments (Ballesteros, 2006, Garrabou *et al.* 2002), the monitoring is recommended to be always performed at the same period of the year in order to facilitate comparisons between years and sites.
44. These sites should be selected according to
- (1) their representativeness at a large geographical scale,
 - (2) their accessibility and
 - (3) the logistical facilities that may contribute to guarantee and facilitate the monitoring operations.

We recommend the setup of sentinel/reference sites in fully protected zones within Marine Protected Areas. MPAs offer excellent facilities for long-term studies and are optimum conditions to approach to the “pristine” functioning of coralligenous assemblages. This precious information would serve as reference for guiding the adoption of conservation and management goals at different international and national levels.

IV.3. FUNCTIONING

45. Special care is to be taken for the study of the functioning of particular associations and species. Specifically, long-lived plants and animals that usually are the engineering species of the coralligenous or the most abundant calcareous algae in maërl beds, need a detailed knowledge of their growth, demographic patterns, vulnerability to disturbances and recovery capacities.
46. Research actions to fill the gaps of current knowledge should focus on
- (a) Bioconcretion dynamics (building and erosion processes);
 - (b) Population dynamics of typical/indicator species;
 - (c) Establish response of key/typical species to different stressors.

V. CONSERVATION ACTIVITIES

V.1. MAJOR THREATS

47. Major threats affecting coralligenous/maërl communities roughly coincide with threats affecting Mediterranean marine biodiversity and are listed in the Strategic Action Programme for the Conservation of Biological Diversity (SAP BIO). However, due to its special habitat and features, not all the threats listed in the SAP BIO affect coralligenous/maërl communities, but some of them are especially relevant. It follows a brief description of the main threats.

V.1.1. TRAWLING

48. Trawling is probably the most destructive impact currently affecting coralligenous communities. Trawling is also completely destructive in maërl beds, being the main cause of maërl disappearance in large Mediterranean areas. The action of trawling gear over coralligenous/maërl assemblages leads to the death of most engineering, dominant and builder species, completely changing the environmental conditions of the coralligenous microhabitats and the maërl environment. As most of these species are particularly long-lived, have low recruitment and complex demographic patterns, destruction of the coralligenous/maërl structure is critical as their recovery will probably take several decades or even centuries. Trawling has also a great impact on target species that, although not as vulnerable as most suspension feeders, they also suffer from this indiscriminate method of fishing.
49. Finally, even the performance of trawling close to coralligenous outcrops or maërl beds affects negatively to algal growth and suspension-feeding due to an increase in turbidity and sedimentation.

V.1.2. ARTISANAL AND RECREATIONAL FISHING

50. Both traditional and recreational fishing also have an effect on coralligenous communities, although they mainly affect the target species. Fishing leads to a significant decrease in mean specific number of fish species, producing changes in the community composition. Certain fishes, mainly elasmobranchs, are severely decimated by artisanal fishing practices when fishing pressure is outstanding. This is the case, for example, of several small sharks such as *Scyliorhinus stellaris*, *Mustelus spp.* or *Squalus spp.* In several places, other species such as groupers and lobsters (e.g. *Epinephelus marginatus* and *Palinurus elephas*) need the implementation of adequate fishery management. Besides, fishing activities can degrade habitat complexity due to the breakage and mortality of fragile macrobenthic species during contact with fishing lines and nets (Bavestrello et al. 2000). The consequent erosion of complexity results from the reduction in the abundance and/or size of large gorgonian and other erect species (e.g. *Axinella spp.*, *Hornera frondiculata*) (Tunesi et al., 1991). The reduction of complexity could infer further biodiversity loss, however the extent of this impact and the associated mechanisms are still poorly understood (Cerrano et al. 2010).
51. Special care has to be taken with the commercial exploitation of red coral (*Corallium rubrum*), whose stocks have strongly declined in most areas. Adequate management of this extremely valuable and long-lived species is necessary.

V.1.3. ANCHORING

52. Anchoring has a very severe impact on coralligenous concretions, as most of the engineering organisms are very fragile and are easily detached or broken by anchors and chains. Coralligenous concretions of frequently visited sites by recreational fishing or diving activities are degraded by the destructive potential of anchors.

V.1.4. INVASIVE SPECIES

53. Currently, at least three algal species are threatening coralligenous/maërl communities in the Western Mediterranean: *Womersleyella setacea*, *Acrothamnion preissii*, *Caulerpa racemosa v. cylindracea* and *Caulerpa taxifolia* (e.g. Cebrian et al. 2012, De Caralt & Cebrian 2013, Cebrian & Rodríguez-Prieto 2012). All of them are only invasive in relatively shallow water coralligenous outcrops and maërl beds (until 60 m), where irradiance levels are sufficient to permit their growth. However, they are especially dangerous, because they completely cover the basal stratum of encrusting corallines and increase sedimentation rates which lead to a total shut down of coralligenous growth or the survival of rhodolites. Most studies have been carried in the Western Mediterranean. There is an absolute lack of knowledge on the effects that lessepsian species have on coralligenous/maërl communities in the Eastern Mediterranean.

V.1.5. GLOBAL WARMING

54. Anomalous high water temperatures were concomitant with large scale mortalities of several suspension feeders (mainly sponges and anthozoans) growing in coralligenous assemblages (Cerrano et al., 2000; Garrabou et al. 2009). Thus, it is expected that under the current trend of global warming (Somot et al. 2008), coralligenous assemblages will be affected by new mortality events during next decades especially in areas where coralligenous assemblages are situated above the summer level of the thermocline.

V.1.6. WASTE WATER DISCHARGES

55. Waste waters profoundly affect the structure of coralligenous communities by inhibiting coralline algal growth, increasing bioerosion rates, decreasing species richness and densities of the largest individuals of the epifauna, eliminating some taxonomical groups and increasing the abundance of highly tolerant species (Hong, 1980, 1982; Cormaci et al., 1985; Ballesteros, 2006). Although no information is available on the impact of eutrophication in Mediterranean maërl beds, the effects must be similar to those reported for coralligenous concretions.

V.1.7. AQUACULTURE

56. Although there are no studies on the impact of aquaculture facilities situated over or at the proximity of coralligenous outcrops, nor maërl beds, their effects should match those produced by waste water dumping.

V.1.8. CHANGES IN LAND USE AND COASTAL INFRASTRUCTURE CONSTRUCTION AND URBANIZATION

57. Most anthropogenic changes in coastal areas or at their vicinity involve an increase in water turbidity and/or sediment removal that affect coralligenous/maërl communities.

V.1.9. RECREATIONAL ACTIVITIES (EXCLUDING FISHING)

58. Uncontrolled or over-frequentation of divers in coralligenous communities has been described to produce an important effect over certain large or fragile suspension feeders inhabiting coralligenous communities (Sala et al., 1996; Garrabou et al., 1998; Coma et al., 2004; Linares et al. 2012).

V.1.10. MUCILAGINOUS AND FILAMENTOUS ALGAL AGGREGATES

59. Blooms of mucilaginous and filamentous algal aggregates can cause severe damage over erect suspension feeders (mainly gorgonians). These blooms are still not well understood but they are apparently caused by eutrophication (Giuliani et al. 2005, Danovaro et al. 2009).

V.2. LEGISLATION AND REGULATIONS

60. Coralligenous/maërl assemblages should be granted legal protection at the same level as *Posidonia oceanica* meadows. A first step would be the inclusion of coralligenous concretions and maërl beds as a priority natural habitat type in the EU Habitats Directive (92/43/EU), which would enable EU countries to undertake surveillance of the conservation status of coralligenous/maërl assemblages and also to set an ecological network of areas of conservation (LICs/ZECs) hosting coralligenous/maërl assemblages, which would ensure their conservation or restoration at a favorable conservation status. Although *Phymatolithon calcareum* and *Lithothamnion corallioides* are present in the Annex V of the Habitats Directive and as such they should be provided by management measures in case of exploitation (which is never the case in the Mediterranean), there is no specific protection for maërl beds. Similar actions should be encouraged in non-EU countries through the existing tools of the Barcelona Convention.
61. Regarding again European countries, recently (21 December 2006), it was published a Council Regulation (EC) No 1967/2006 concerning management measures for the sustainable exploitation of fishery resources in the Mediterranean Sea, amending Regulation (EU) No 2847/93 and repealing Regulation (EC) No 1626/94 which states that "Fishing trawl nets, dredges, shore seines or similar nets above coralligenous habitats and maërl beds shall be prohibited" (Article 4.2) and that this prohibition "shall apply to all Natura 2000 sites, all special protected areas and all specially protected areas of Mediterranean importance (SPAMIs) which have been designated for the purpose of the conservation of these habitats under either Directive 92/43/EU or Decision 1999/800/EU" (Article 4.4).
62. In 2008 the European Union adopted the Marine Strategy Framework Directive (MSFD 2008/56/EC) which requires to maintain European marine waters in "Good Environmental Status" (GES). The MSFD included 11 descriptors for the assessment of GES among them the Sea-floor Integrity is defined as "Sea-floor integrity is at a level that ensures that the structure and functions of the ecosystems are safeguarded and benthic ecosystems, in particular, are not adversely affected." (Rice et al. 2012). This descriptor directly concerns biogenic structures such as the Mediterranean

coralligenous and different initiatives are underway to determine the GES of coralligenous habitats (e.g. Gatti et al. 2015). The monitoring of different indicators (such as those indicated in this document and other proposed by other authors) should allow determining reference conditions at regional scales and the proposal of a quantitative index to evaluate the GES in each area. The final aim of MSFD is to guide management and conservation actions for maintaining and when necessary recovering the good environmental status of waters.

63. In line with the MSFD, the contracting parties to the Barcelona Convention set targets for achieving GES of the Mediterranean Sea and its coastal zone by 2020. In achieving these targets it has been recognized the importance to apply the ecosystem approach (EcAp) to the management of human activities that may affect the Mediterranean marine and coastal environment for the promotion of sustainable development (UNEP/MAP 2007). In addition, through Decision IG 21/3 (the so-called “COP 18 EcAp Decision”) the contracting Parties agreed to design an Integrated Monitoring and Assessment Program for the next meeting of the contracting parties (COP19) and mandated the Secretariat to carry out an assessment of the state of the Mediterranean environment in 2017 which necessarily will include the coralligenous/maërl habitats (UNEP/MAP, 2013).

V.3. CREATION OF MARINE PROTECTED AREAS

64. Within the Convention on Biological Diversity (CBD) countries have committed to protect by 2020 “10 % of coastal and marine areas, especially areas of particular importance for biodiversity and ecosystem services, are conserved through effectively and equitably managed, ecologically representative and well-connected systems of protected areas and the effective area-based conservation measures” (Target 11 of the Aichi Strategic Plan for Biodiversity 2020) and the Roadmap for a comprehensive coherent network of well managed MPAs to achieve Aichi Target 11 in the Mediterranean. Overall, only about 1 % of Mediterranean coastal waters susceptible to harbor coralligenous/maërl assemblages are protected.
65. Most present Mediterranean MPAs are devoted to protect *Posidonia oceanica* meadows and other shallow water assemblages, in such a way that the percentage of coralligenous/maërl habitat currently protected in the Mediterranean is extremely low. Thus, it is necessary to protect representative coralligenous/maërl assemblages by applying the protection and management measures recommended by Articles 6 and 7 of the SPA/BD protocol. In fact, Marine Protected Areas (MPAs) have to be established taking into account the seascape diversity and trying to include places with several relevant assemblages, as has been already applied in the creation and zonation of some MPAs (Villa *et al.*, 2002; Di Nora *et al.*, 2007).
66. Countries have to identify and cartography as soon as possible sea bottoms covered by coralligenous outcrops and maërl beds in order to design a network of MPAs that enables the protection of coralligenous/maërl assemblages.
67. Those Mediterranean MPAs, which contain coralligenous/maërl assemblages and for which management and monitoring plans have not yet been developed and implemented, must develop and implement such plans as soon as possible.

V.4. NATIONAL PLANS

68. To ensure more efficiency in the measures envisaged in the implementation of this Action Plan, Mediterranean countries are invited to establish national plans for the conservation of Coralligenous and other calcareous bio-concretions. Each national plan should take into account the concerned country's, or even areas', specific features. It must suggest appropriate legislative measures, particularly for the environmental impact assessment of coastal infrastructure (building works, pipelines out to sea, and deposits of material from dredging) and to control activities which could affect coralligenous/maërl assemblages. The national plan shall be based on the available scientific data and will include programmes for
- (i) collection and regular updating of data,
 - (ii) training and refresher courses for specialists,
 - (iii) awareness-raising and education for the general public, actors and decision-makers and
 - (iv) the conservation of coralligenous/maërl assemblages of importance for the Mediterranean marine environment.

The national plans must be brought to the attention of all concerned actors and, when possible, coordinated with the relevant national plans (e.g. emergency plan to deal with pollution).

VI. COORDINATION OF THIS ACTION PLAN WITH OTHER TOOLS AND INITIATIVES

69. The Standard Data-Entry Form (SDF), developed by SPA/RAC, can be used to identify potentially good sites for the establishment of MPAs devoted to protect coralligenous/maërl assemblages. Besides the analysis of current data on the distribution of coralligenous assemblages along with information derived from distribution modelling tools can help guiding cost-effective future surveys and monitoring efforts towards the development of basin-wide marine protected areas network for coralligenous/maërl assemblages (Martin et al. 2014).
70. However the SDF is not appropriate to be used in the monitoring of coralligenous/maërl assemblages since it has been designed for the inventory of sites and habitats but not for an accurate assessment of multi-species population densities and their evolution. Annex B (habitat types) from the SDF should be slightly modified in the point IV.3.1 (Coralligenous biocenosis), according to current knowledge. Species appearing in Annex C should be slightly enlarged in order to include several engineering coralligenous species according to the adopted criteria for amendments of the Annexes (II & III) of the Protocol SPA/BD.
71. MPAs classified as SPAMIs and containing coralligenous/maërl assemblages inside their protected areas should develop management and protection plans to ensure their conservation.

VII. REGIONAL COORDINATION STRUCTURE

72. Regional coordination of the implementing of the present Action Plan will be guaranteed by the Mediterranean Action Plan's (MAP) secretariat through the Specially Protected Areas Regional Activity Centre. The main functions of the coordinating structure shall consist in:
- collecting, validating and circulating data at Mediterranean level;
 - promoting the drawing up of inventories of species, coralligenous/maërl assemblages of importance for the Mediterranean marine environment;
 - promoting trans-boundary cooperation;
 - promoting and supporting the setting up of coralligenous/maërl assemblages monitoring networks;
 - preparation of reports on progress in the implementation of the Action Plan, to be submitted to the meeting of national focal points for SPAs and to meetings of the Contracting Parties;
 - organizing meetings of experts on specific subjects relating to coralligenous/maërl assemblages and training sessions.
73. Complementary work done by other international organizations, and aiming at the same objectives, shall be encouraged, promoting coordination and avoiding possible duplication of efforts.

VIII. PARTICIPATION IN THE IMPLEMENTATION

74. Implementing the present Action Plan is the province of the national authorities of the Contracting Parties. The concerned international organizations and/or NGOs, laboratories and any organization or body are invited to join in the work necessary for implementing the present Action Plan. At their ordinary meetings, the Contracting Parties may, at the suggestion of the meeting of National Focal Points for SPAs, grant the status of «Action Plan Associate» to any organization or laboratory which so requests and which carries out, or supports (financially or otherwise) the carrying out of concrete actions (conservation, research, etc.) likely to facilitate the implementation of the present Action Plan, taking into account the priorities contained therein.
75. The coordination structure shall set up a mechanism for regular dialogue between the participating organizations and, where necessary, organize meetings to this effect. Dialogue should be made mainly by mail, including E-mail.

ANNEX: IMPLEMENTATION TIMETABLE


Action	Deadline	To be implemented by
1. Build and publish the database of scientists and research institutions working on the coralligenous assemblages and maërl beds.	2016	SPA/RAC
2. Guidelines for the assessment of environmental impact on coralligenous/maërl assemblages	2017	SPA/RAC
3. Development of Working Groups on coralligenous assemblages and maërl beds.	2016	SPA/RAC-Contracting Parties
4. Build-up a coralligenous/maërl assemblages distribution online database	2018	SPA/RAC-Contracting Parties
5. Improve habitat modeling methods could provide new predictive models on coralligenous distribution and guide cost-effective field surveys for data acquisition	2017	Contracting Parties
6. Characterization of coralligenous habitats at regional scale	2018	SPA/RAC-Contracting Parties
7. Build-up a Check-list / Reference species list for the coralligenous assemblages	2016	SPA/RAC
8. Development of standardized protocols for the characterization of coralligenous / maërl assemblages.	2017	SPA/RAC-Contracting Parties
9. Development of indices and/or intercalibration initiatives to determine conservation environmental status of coralligenous	2017	SPA/RAC-Contracting Parties
10. Set a network of sentinel sites on coralligenous across the Mediterranean	2020	SPA/RAC-Contracting Parties
11. Promote research programs on coralligenous assemblages and maerl beds	2016	Contracting Parties
12. Develop and implement legislation initiatives for the conservation of coralligenous assemblages	Ongoing	Contracting Parties
13. Coordinate the design of an Integrated Monitoring and Assessment Program for the assessment of the state coralligenous/maërl assemblages in view to be included the assessment of the state of the Mediterranean	2016	Contracting Parties
14. Promote the declaration of marine protected areas to preserve coralligenous assemblages in coastal and offshore areas	2018	SPA/RAC-Contracting Parties
15. Build-up a coordination platform on different initiatives devoted to the coralligenous/maërl assemblages	2017	SPA/RAC
16. Organize a Symposium on coralligenous assemblages and maërl beds every 3 years	2018	SPA/RAC
17. Preparation of a communication plan to raise the awareness on the importance of coralligenous assemblages and maërl beds for the conservation of Mediterranean biodiversity	2017	SPA/RAC

IX. REFERENCES

- Ballesteros, E. 1991. Seasonality of growth and production of a deep-water population of *Halimeda tuna* (Chlorophyceae, Caulerpales) in the North-western Mediterranean. *Bot. Mar.* 34: 291-301.
- Ballesteros, E. 2006. Mediterranean coralligenous peuplements: a synthesis of present knowledge. *Oceanogr. Mar. Biol. Ann. Rev.* 44: 123-195.
- Belsher, T., Houlgatte, E., Boudouresque, C.F. 2005. Cartographie de la prairie à *Posidonia oceanica* et des principaux faciès sédimentaires marins du Parc National de Port-Cros (Var, France, Méditerranée). *Sci. Rep. Port-Cros nat. Park* 21: 19-28.
- Bianchi, C.N., Pronzato, R., Cattaneo-Vietti, R., Benedetti-Cecchi, L., Morri, C., Pansini, M., Chemello, R., Milazzo, M., Fraschetti, S., Terlizzi, A., Peirano, A., Salvati, E., Benzoni, F., Calcinai, B., Cerrano, C., Bavestrello, G. 2004. Hard bottoms. *Biol. Mar. Medit.* 11 (suppl. 1): 185-215.
- BIOMAERL Team, 2003. Conservation and management of Northeast Atlantic and Mediterranean Maerl Beds. *Aquatic Conservation. Marine and Freshwater Ecosystems*, 13 (suppl. 1): 65-76.
- Boudouresque, C. F. 1971. Méthodes d'étude qualitative et quantitative du benthos (en particulier du phytobenthos). *Téthys* 3: 79-104.
- Braun-Blanquet, J. 1979. Fitosociología. Blume. Madrid.
- Casas, E., Teixidó, N., Garrabou, J., Cebrian, E. 2015. Structure and biodiversity of coralligenous peuplements over broad spatial and temporal scales. *Mar. Biol.* 162:901–912
- Cebrian, E., Rodríguez-Prieto, C., 2012. Marine Invasion in the Mediterranean Sea: The Role of Abiotic Factors When There Is No Biological Resistance. *PLoS ONE* 7(2): e31135. doi:10.1371/journal.pone.0031135
- Cebrian, E., Linares, C., Marschal, C., Garrabou, J. 2012. Exploring the effects of invasive algae on the persistence of gorgonian populations. *Biol. Inv.* 14: 2647–2656 DOI: 10.1007/s10530-012-0261-66
- Cebrian, E., Ballesteros, E. 2004. Zonation patterns of benthic communities in an upwelling area from the western Mediterranean (La Herradura, Alboran Sea). *Sci. Mar.* 68: 69-84.
- Cecchi, E., Piazzoli, L. 2010. A new method for the assessment of the ecological status of coralligenous assemblages. *Biol. Mar. Mediterr.* 17(1), 162–163.
- Cerrano, C., Danovaro, R., Gambi, C., Pusceddu, A., Riva, A., Schiaparelli S (2010) Gold coral (*Savalia savaglia*) and gorgonian forests enhance benthic biodiversity and ecosystem functioning in the mesophotic zone. *Biodivers. Conserv.* 19:153-167.
- Cerrano, C., Bavestrello, G., Bianchi, C.N., Cattaneo-Vietti, R., Bava, S., Morganti, C., Morri, C., Picco, P., Sara, G., Schiaparelli, S., Siccardi, A., Sponga, F. 2000. A catastrophic mass-mortality episode of gorgonians and other organisms in the Ligurian Sea (NW Mediterranean), summer 1999. *Ecol. Lett.* 3: 284-293.
- Coma, R., Linares, C., Ribes, M., Díaz, D., Garrabou, J., Ballesteros, E. 2006. Consequences of a mass mortality in populations of *Eunicella singularis* (Cnidaria: Octocorallia) in Menorca (NW Mediterranean). *Mar. Ecol. Progr. Ser.* 327: 51-60.
- Coma, R., Polà, E., Ribes, M., Zabala, M. 2004. Long-term assessment of temperate octocoral mortality patterns, protected vs. unprotected areas. *Ecol. Appl.* 14: 1466-1478.
- Cormaci, M., Furnari, G., Giaccone, G. 2004. Macrophytobenthos. *Biol. Mar. Medit.* 11(suppl. 1): 217-246.
- Cormaci, M., Furnari, G., Scamacca, B. 1985. Osservazioni sulle fitocenosi bentoniche del golfo di Augusta (Siracusa). *Bollettino dell'Accademia Gioenia Scienze Naturali* 18: 851-872.
- Dailianis, T., Tsigenopoulos, C.S., Dounas, C., Voultsiadou, E. 2014. Genetic diversity of the imperilled bath sponge *Spongia officinalis* Linnaeus, 1759 across the Mediterranean Sea: patterns of population differentiation and implications for taxonomy and conservation. *Molec. Ecol.* 20:3757-3772
- Danovaro, R., Fonda Umani, S., Pusceddu, A. 2009. Climate Change and the potential spreading of marine mucilage and microbial pathogens in the Mediterranean Sea. *PLoS ONE* 4(9): e7006
- De Caralt, S., Cebrian, E. 2013. Impact of an invasive alga (*Womersleyella setacea*) on sponge assemblages: compromising the viability of future populations. *Biol. inv.* 15:1591-1608
- Deter, J., Descamp, P., Ballesta, L., Boissery, P., Holon, F. 2012. A preliminary study toward an index based on coralligenous assemblages for the ecological status assessment of Mediterranean French coastal waters. *Ecol. Indicat.* 20:345–352.

- Di Nora, T., Agnesi, S., Tunesi, L. 2007. Planning of marine protected areas: useful elements to identify the most relevant scuba-diving sites. *Rapp. Comm. int. Mer Médit.*, 38.
- Fraschetti, S., Bianchi, C.N., Terlizzi, A., Fanelli, G., Morri, C., Boero, F. 2001. Spatial variability and human disturbance in shallow subtidal hard substrate assemblages: a regional approach. *Mar. Ecol. Progr. Ser.* 212: 1-12.
- García-Carrascosa, A.M. 1987. El bentos de los alrededores de las Islas Columbretes. Elementos para su cartografía bentónica. In: *Islas Columbretes: Contribución al estudio de su medio natural*. L. A. Alonso, J.L. Carretero & A.M. García-Carrascosa (coords.). COPUT, Generalitat Valenciana, Valencia: 477-507.
- García-Rubies, A, Hereu, B, Zabala, M. 2013. Long-Term Recovery Patterns and Limited Spillover of Large Predatory Fish in a Mediterranean MPA. *PLoS ONE* 8(9): e73922. doi:10.1371/journal.pone.0073922
- Garrabou, J., Kipson, S., Kaleb, S., Kruzic, P., Jaklin, A., Zuljevic, A., Rajkovic, Z., Rodic P., Jelic, K., Zupan, D. 2014. Monitoring Protocol for Reefs - Coralligenous Community, MedMPAnet Project
- Garrabou, J., Coma, R., Bally, M., Bensoussan, N., Chevaldonné, P., Cigliano, M., Diaz, D., Harmelin, J.G., Gambi, M.C., Kersting, D.K., Lejeusne, C., Linares, C., Marschal, C., Pérez, T., Ribes, M., Romano, J.C., Serrano, E., Teixido, N., Torrents, O., Zabala, M., Zuberer, F., Cerrano, C. 2009. Mass mortality in northwestern Mediterranean rocky benthic communities : effects of the 2003 heat wave. *Global Change Biology* 15:1090-1103
- Garrabou, J. 1998. Applying a Geographical Information System (GIS) to the study of growth of benthic clonal organisms. *Mar. Ecol. Progr. Ser.* 173: 227-235.
- Garrabou, J. 1999. Life history traits of *Alcyonium acaule* and *Parazoanthus axinellae* (Cnidaria, Anthozoa), with emphasis on growth. *Mar. Ecol. Progr. Ser.* 178: 193-204.
- Garrabou, J., Ballesteros, E. 2000. Growth of *Mesophyllum alternans* and *Lithophyllum frondosum* (Corallinaceae, Rhodophyta) in the Northwestern Mediterranean. *Eur. J. Phycol.* 35: 1-10.
- Garrabou, J., Ballesteros, E., Zabala, M. 2002. Structure and dynamics of north-western Mediterranean rocky benthic communities along a depth gradient. *Est. Coast. Shelf Sci.* 55: 493-508.
- Garrabou, J., Perez, T., Sartoretto, S., Harmelin, J.G. 2001. Mass mortality event in red coral (*Corallium rubrum*, Cnidaria, Anthozoa, Octocorallia) population in the Provence region (France, NW Mediterranean). *Mar. Ecol. Progr. Ser.* 217: 263-272.
- Garrabou, J., Sala, E., Arcas, A., Zabala, M. 1998. The impact of diving on rocky sublittoral communities: a case study of a bryozoan population. *Conserv. Biol.* 12: 302-312.
- Garrabou, J., Zabala, M. 2001. Growth dynamics in four Mediterranean demosponges. *Estuar. Coast. Shelf Sci.* 52: 293-303.
- Gatti G, Bianchi CN, Morri C, Montefalcone M, Sartoretto S. 2015. Coralligenous reefs state a long anthropized coasts: Application and validation of the COARSE index, based on a rapid visual assessment (RVA) approach. *Ecol. Indicat.* 52:567-576
- Gatti, G., Montefalcone, M., Rovere, A., Parravicini, V., Morri, C., Albertelli, G., Bianchi, C.N. 2012. Seafloor integrity down the harbor waterfront: the coralligenous shoals off Vado Ligure (NW Mediterranean). *Adv Ocean Limnol* 3(1):51–67.
- Germonpre, P. 2006. The medical risks of underwater diving and their control. *Int. Sport. J.* 7: 1-15.
- Giakoumi, S. et al. 2013. Ecoregion-Based Conservation Planning in the Mediterranean: Dealing with Large-Scale Heterogeneity. *PLoS One* 8, e76449 (2013).
- Gili, J.M., Ros, J. 1987. Study and cartography of the benthic communities of Medes Islands (NE Spain). *P.S.Z.N.I. Mar. Ecol.* 6: 219-238.
- Harmelin, J.G., Marinopoulos, J. 1994. Population structure and partial mortality of the gorgonian *Paramuricea clavata* (Risso) in the north-western Mediterranean (France, Port-Cros Island). *Marine Life* 4: 5-13.
- Hong, J.S. 1980. Étude faunistique d'un fond de concrétionnement de type coralligène soumis à un gradient de pollution en Méditerranée nord-occidentale (Golfe de Fos). *Thèse de Doctorat. Université d'Aix-Marseille II*.
- Hong, J.S. 1982. Contribution à l'étude des assemblages d'un fond coralligène dans la région marseillaise en Méditerranée Nord-Occidentale. *Bulletin of Korea Ocean Research and Development Institute* 4: 27-51.
- Kipson, S, Fourt, M, Teixido, N, Cebrian, E, Casas, E, Ballesteros, E, Zabala, M, Garrabou, J. 2011. Rapid Biodiversity Assessment and Monitoring Method for Highly Diverse Benthic Communities: a Case Study of Mediterranean Coralligenous Outcrops. *PLoS ONE* 6(11): e27103 doi:10.1371/journal.pone.0027103

- Laborel, J. 1987. Marine biogenic constructions in the Mediterranean. *Sci. Rep. Port-Cros Natl. Park* 13: 97-126.
- Linares, C., Garrabou, J., Hereu, B., Díaz, D., Marschal, C., Sala, E., Zabala, M. 2012. Beyond fishes: assessing the effectiveness of marine reserves on overexploited long-lived sessile invertebrates. *Conserv. Biol.* 26:88-96
- Linares, C. 2006. Population ecology and conservation of a long-lived marine species: the red gorgonian *Paramuricea clavata*. Tesi Doctoral. Universitat de Barcelona. 210 pp.
- Linares, C., Coma, R., Díaz, D., Zabala, M., Hereu, B., Dantart, L. 2005. Immediate and delayed effects of mass mortality event on gorgonian population dynamics and benthic community structure in the NW Mediterranean. *Mar. Ecol. Progr. Ser.* 305: 127-137.
- Linares, C., Doak, D.F., Coma, R., Díaz, D., Zabala, M. *in press*. Life history and population viability of a long-lived marine invertebrate: the octocoral *Paramuricea clavata*. *Ecology*.
- Martin et al. 2014. Coralligenous and maërl habitats: predictive modelling to identify their spatial distributions across the Mediterranean Sea. *Scientific Reports* 4: 5073
- Pérez, J., Picard, J.M. 1964. Nouveau manuel de bionomie benthique de la mer Méditerranée. *Recueil Travaux Station Marine Endoume* 31(47): 1-131.
- Pérez, T., Garrabou, J., Sartoretto, S., Harmelin, J.G., Francour, P., Vacelet, J. 2000. Mortalité massive d'invertébrés marins: un événement sans précédent en Méditerranée nord-occidentale. *Comptes Rendus Académie des Sciences Série III, Life Sciences* 323: 853-865.
- Ramos, A.A. 1985. Contribución al conocimiento de las biocenosis bentónicas litorales de la Isla Plana o Nueva Tabarca (Alicante). In: *La reserva marina de la Isla Plana o Nueva Tabarca (Alicante)*. A.A. Ramos (ed.), Ayuntamiento de Alicante-Universidad de Alicante: 111-147.
- Sala, E., Ballesteros, E. 1997. Partitioning of space and food resources by three fishes of the genus *Diplodus* (Sparidae) in a Mediterranean rocky infralittoral ecosystem. *Mar. Ecol. Progr. Ser.* 152: 273-283.
- Sala, E., Garrabou, J., Zabala, M. 1996. Effects of diver frequentation on Mediterranean sublittoral populations of the bryozoan *Pentapora fascialis*. *Mar. Biol.* 126: 451-459.
- Teixido N, Casas E, Cebrián E, Linares C, Garrabou J (2013). Impacts on coralligenous outcrop biodiversity of a dramatic coastal storm. *PLoS ONE* 10.1371/journal.pone.0053742
- Teixido, N, Garrabou, J Harmelin, J.G. 2011. Low dynamics, high longevity and persistence of sessile structural species dwelling on Mediterranean coralligenous outcrops. *PLoS ONE* 6(8): e23744. doi:10.1371/journal.pone.0023744
- Templado, J., Calvo, M. (eds.). 2002. Flora y Fauna de la Reserva Marina de las Islas Columbretes. Secretaría Gral. De Pesca Marítima, Mº de Agricultura, Pesca y Alimentación, Madrid, 263 pp.
- Templado, J., Calvo, M. (eds.). 2006. Flora y Fauna de la Reserva Marina y Reserva de Pesca de la Isla de Alborán. Secretaría Gral. De Pesca Marítima, Mº de Agricultura, Pesca y Alimentación, Madrid, 269 pp.
- Tetzaff, K., Thorsen, E. 2005. Breathing at depth: physiological and clinical aspects of diving when breathing compressed air. *Clin. Chest Med.* 26: 355-380.
- Trygonis, V., Sini, M., 2012. photoQuad: a dedicated seabed image processing software, and a comparative error analysis of four photoquadrat methods. *J. Exp. Mar. Biol. Ecol.* 424-425, 99-108. doi:10.1016/j.jembe.2012.04.018
- Tunesi, L., Peirano, A., Romeo, G, Sassarini, M., 1991. Problématiques de la protection des faciès à Gorgonaires sur les fonds côtiers de "Cinque Terre" (Mer Ligure, Italie). In: *Les Espèces marines à protéger en Méditerranée* (C.F. Boudouresque, M. Avon & V. Gravez, eds.). GIS Posidonie, Marseille: 65-70.
- UNEP-MAP-SPA/RAC (2011) Draft Lists of coralligenous/ maërl populations and of main species to be considered by the inventory and monitoring. Expert Meeting to propose standard methodologies for the inventory and monitoring of coralligenous/maërl communities and their main species. Rome, Italy, 7-8 April 2011, 11 pp.
- Villa, F., Tunesi, L., Agardy, T. 2002. Optimal zoning of marine protected areas through spatial multiple criteria analysis: the case of Asinara Island National Marine Reserve of Italy. *Conserv. Biol.* 16: 1-12.



United Nations
Environment Programme



Mediterranean Action Plan
Barcelona Convention

United Nations Environment Programme
Coordinating Unit for the Mediterranean Action Plan
Barcelona Convention Secretariat
48, Vassileos Konstantinou Avenue 11635
P.O. Box: 18019, Athens Greece
Tel: +302107273100 - Fax: +302107253196
www.unepmap.org